

2016 WHO Medical Eligibility Criteria for Contraceptive Use: Quick Reference Chart for Category 3 and 4

to initiate or continue use of combined oral contraceptives (COCs), depot-medroxyprogesterone acetate (DMPA), progestin-only implants, copper intrauterine device (Cu-IUD), levonorgestral intrauterine system (LNG-IUS)

CONDITION	Sub-condition	COC	DMPA	Implants	Cu-IUD	LNG-IUS
Pregnancy		NA	NA	NA		
Breastfeeding	Less than 6 weeks postpartum					
	≥ 6 weeks to < 6 months postpartum				See i.	See i.
	≥ 6 months postpartum					
Postpartum not breastfeeding VTE = venous thromboembolism	< 21 days					
	< 21 days with other risk factors for VTE*				See i.	See i.
	≥ 21 to 42 days with other risk factors for VTE*					
Postpartum timing of insertion	≥ 48 hours to less than 4 weeks					
	Puerperal sepsis	See i.	See i.	See i.		
Postabortion (immediate post-septic)						
Smoking	Age ≥ 35 years, < 15 cigarettes/day					
	Age ≥ 35 years, ≥ 15 cigarettes/day					
Multiple risk factors for cardiovascular disease						
Hypertension BP = blood pressure	History of (where BP cannot be evaluated)					
	BP is controlled and can be evaluated					
	Elevated BP (systolic 140-159 or diastolic 90-99)					
	Elevated BP (systolic ≥ 160 or diastolic ≥ 100)					
	Vascular disease					
Deep venous thrombosis (DVT) and pulmonary embolism (PE)	History of DVT/PE					
	Acute DVT/PE					
	DVT/PE, established on anticoagulant therapy					
	Major surgery with prolonged immobilization					
Known thrombogenic mutations						
Ischemic heart disease (current or history of)				I C		I C
Stroke (history of)				I C		
Complicated valvular heart disease						
Systemic lupus erythematosus	Positive or unknown antiphospholipid antibodies					
	Severe thrombocytopenia		I C		I C	

Adapted from: *Medical Eligibility Criteria for Contraceptive Use, 5th Edition*. Geneva: World Health Organization, 2015. Available: http://www.who.int/reproductivehealth/publications/family_planning/en/index.html

- Category 1** There are no restrictions for use.
- Category 2** Generally use; some follow-up may be needed.
- Category 3** Usually not recommended; clinical judgment and continuing access to clinical services are required for use.
- Category 4** The method should not be used.

CONDITION	Sub-condition	COC	DMPA	Implants	Cu-IUD	LNG-IUS
Headaches	Migraine without aura (age < 35 years)	I C				
	Migraine without aura (age ≥ 35 years)	I C				
	Migraines with aura (at any age)		I C	I C		I C
Unexplained vaginal bleeding (prior to evaluation)						
Gestational trophoblastic disease	Regressing or undetectable β-hCG levels					
	Persistently elevated β-hCG levels or malignant disease					
Cancers	Cervical (awaiting treatment)				I C	I C
	Endometrial				I C	I C
	Ovarian				I C	I C
Breast disease	Current cancer					
	Past w/ no evidence of current disease for 5 yrs					
Uterine distortion (due to fibroids or anatomical abnormalities)						
STIs/PID	Current purulent cervicitis, chlamydia, gonorrhea				I C	I C
	Current pelvic inflammatory disease (PID)				I C	I C
	Very high individual risk of exposure to STIs				I C	I C
Pelvic tuberculosis						
Diabetes	Nephropathy/retinopathy/neuropathy					
	Diabetes for > 20 years					
Symptomatic gall bladder disease (current or medically treated)						
Cholestasis (history of related to oral contraceptives)						
Hepatitis (acute or flare)						
Cirrhosis (severe)						
Liver tumors (hepatocellular adenoma and malignant hepatoma)						
AIDS	No antiretroviral (ARV) therapy	See ii.	See ii.	See ii.	I C	I C
	Not improved on ARV therapy				I C	I C
Drug interactions	Rifampicin or rifabutin					
	Anticonvulsant therapy**					

This chart shows a complete list of all conditions classified by WHO as Category 3 and 4. Characteristics, conditions, and/or timing that are Category 1 or 2 for all methods are not included in this chart (e.g., menarche to < 18 years, being nulliparous, obesity, high risk of HIV or HIV-infected, < 48 hours and more than 4 weeks postpartum).

- I/C** Initiation/Continuation: A woman may fall into either one category or another, depending on whether she is initiating or continuing to use a method. Where I/C is not marked, the category is the same for initiation and continuation.
- NA** Not Applicable: Women who are pregnant do not require contraception. If these methods are accidentally initiated, no harm will result.
- i** The condition, characteristic and/or timing is not applicable for determining eligibility for the method.
- ii** Women who use methods other than IUDs can use them regardless of HIV/AIDS-related illness or use of ART.
- *** Other risk factors for VTE include: previous VTE, thrombophilia, immobility, transfusion at delivery, BMI > 30 kg/m², postpartum hemorrhage, immediately post-caesarean delivery, pre-eclampsia, and smoking.
- **** Anticonvulsants include: phenytoin, carbamazepine, barbiturates, primidone, topiramate, oxcarbazepine, and lamotrigine. Lamotrigine is a category 1 for implants.